

# A conjecture on square roots of Laplacian and signless Laplacian eigenvalues of graphs

S. Akbari<sup>a,b</sup>, E. Ghorbani<sup>a,b</sup>, M. R. Oboudi<sup>a</sup>

<sup>a</sup>*Department of Mathematical Sciences, Sharif University of Technology, Tehran, Iran*

<sup>b</sup>*School of Mathematics, Institute for Research in Fundamental Sciences (IPM),*

*P.O. Box 19395-5746, Tehran, Iran*

Let  $G$  be a graph with  $n$  vertices,  $m$  edges, the incidence matrix  $X$ , and the directed incidence matrix  $D$  for a given orientation.  $X$  and  $D$  are  $n$ -by- $m$  matrices,  $X$  is a  $(0,1)$ -matrix, and  $D$  a  $(-1,0,1)$ -matrix. Moreover

$$XX^\top = \Delta + A, \text{ and } DD^\top = \Delta - A,$$

where  $A$  is the adjacency matrix and  $\Delta$  is the diagonal matrix whose diagonal entries are the vertex degrees of  $G$ . Note that  $\Delta - A$  is the Laplacian matrix and  $\Delta + A$  is the signless Laplacian matrix of  $G$ . Thus the singular values of  $X$  and  $D$  are the square roots of the signless Laplacian and the Laplacian eigenvalues of  $G$ , respectively. We denote the sum of all singular values of a matrix  $M$  by  $\mathcal{E}(M)$ . We have the following conjecture:

**Conjecture.** For any graph,  $\mathcal{E}(D) \leq \mathcal{E}(X)$ .

It is not hard to see that for bipartite graphs,  $\mathcal{E}(D) = \mathcal{E}(X)$ . A computer aid search showed that the conjecture is true for all graphs up to 9 vertices. For  $k$ -regular graphs with adjacency eigenvalues  $\lambda_1, \dots, \lambda_n$  the conjecture is equivalent to the following inequality:

$$\sqrt{k - \lambda_1} + \dots + \sqrt{k - \lambda_n} \leq \sqrt{k + \lambda_1} + \dots + \sqrt{k + \lambda_n}.$$